

ECC966 INTERGARD 740 MID BLUE PART A

4

10/25/14

1.

1.1. INTERGARD 740 MID BLUE PART A  
ECC966

1.2.

1.3.

626-6

( 8-6 )

1.4.

055-632-6286( ),055 586 2310( )

055 587 6276( )

055 586 2310( )

055 586 2310( )

2.

2.1.

3; H226

/

2;H315

/

2;H319

1;H317

-1 ;H335

3;H412

2.2.

11 , 12



H226  
 H315  
 H317  
 H319  
 H335  
 H412

[ ]:

P210 / / /  
 P260 / /  
 P261 / /가 / / /  
 P262 , ,  
 P264  
 P271 가  
 P272  
 P273  
 P280 / / /

[ ]:

P301+310 : /  
 P302+352 :  
 P303+361+353 ( ) :  
 /  
 P304+312 : /  
 P305+351+338 가 : .가  
 P312 /  
 P321 ( ).  
 P331  
 P333+313 /  
 P337 :  
 P340 가  
 P362  
 P363  
 P370 :  
 P378 , , ,

[ ]:

P403+233 가  
 P405 가

[ ]:

P501 ( )

2.3. PBT ( , ) vPvB ( , )

3.

/	%	GHS	
xylene CAS No: 0001330-20-7	20-30	3; H226 - 4;H312 - 4;H332 / 2;H315	[1][2]

		/ 2AIH319 -1 ;H336 - 1;H372	
Epoxy resin (av.mol.wt.<700) CAS No: 0025068-38-6	10-20	/ 2;H319 / 2;H315 1;H317 - 2;H411	[1]
Cyclohexanol, 4,4(1-methylethylidene)bis-, polymer with (chloromethyl)oxirane CAS No: 0030583-72-3	5-10	1;H317 - 2;H411	[1]
Titanium dioxide CAS No: 0013463-67-7	2.5-5		[1][2]
Ethylbenzene CAS No: 0000100-41-4	2.5-5	2;H225 - 4;H332 - 3;H373 1;H304 / 2;H315 / 2;H319 -1 ;H335	[1][2]
Propylene glycol mono methyl ether CAS No: 0000107-98-2	2.5-5	3; H226 -1 ;H336	[1][2]
Solvent naphtha (petroleum), light aromatic CAS No: 0064742-95-6	1-2.5	1;H304	[1]
1-Methoxy-2-propyl acetate CAS No: 0000108-65-6	1-2.5	3; H226	[1]
polyamide dispersion CAS No: 0055349-01-4	<1	1;H317 - 4;H413	[1]
	40-50	---	---

- 1)  
2) 가  
3) PBT vPvB  
16

4.

4.1.

가

가

4.2. 가 /

4.3.

5.

5.1.

; , , , .  
.

Note; 가 .

가 .

5.2.

, , , 가 :  
.

5.3.

가 .  
가 .

6.

6.1.

가 가 가 , 가

6.2.

가 .

6.3.

.8 .

, , 가 . (13 .)

가 .

가 .

, 가 , 가 . ,  
가 .

7.

7.1.

가 .

가 (LEL) (OEL)

가

가

가 (LEL) (OEL)

7.2. ( )

, 8

가

가

61

가

1

7.3. Specific end use(s)

가

, 가

. 3

Hot surfaces, Sparks,

가

( 60% , )

8.

8.1.

(OEL)

(ACGIH)

(ACGIH)

ppm

mg/m<sup>3</sup>

ppm

mg/m<sup>3</sup>

Barium Sulphate

2

10

Ethylbenzene

125

545

100

435

Propylene glycol mono methyl ether

150

540

100

360

Talc				2
Titanium dioxide				10
xylene	150	655	100	434

(P) (Peak exposure limit)  
(R)  
(Sk)  
(Sen)  
(Cat 1)  
(Cat 2) 가  
(Cat 3)

**DNEL/PNEC**

**8.2.**

가

가

(visor)

(overall)

가

가

가

가

pH

/ (°C)

(°C)

117

27

( = 1)

( , )

/

: 1.1 ( xylene )

: 6.6 ( xylene )

(Pa)

1.32

n-

/

(Log Kow)

### 9.2.

## 10.

### 10.1.

### 10.2.

.(Section 7 )

가

### 10.3.

가

### 10.4.

.(7 .)

### 10.5.

### 10.6.

가

## 11.

(OEL)

가

가

Data

	LD50, mg/kg	LD50, mg/kg	LD50, mg/L/4hr	/ LD50, mg/L/4hr
1-Methoxy-2-propyl acetate - (108-65-6)	8,532.00,	5,000.00,		
Cyclohexanol, 4,4(1-methylethylidene) bis-, polymer with (chloromethyl)oxirane - (30583-72-3)				
Epoxy resin (av.mol.wt.<700) - (25068-38-6)	2,000.00,	2,000.00,		
Ethylbenzene - (100-41-4)	3,500.00,	15,433.00,	17.20,	
polyamide dispersion - (55349-01-4)				
Propylene glycol mono methyl ether - (107-98-2)	5,000.00,	13,000.00,		
Solvent naphtha (petroleum), light aromatic - (64742-95-6)	6,800.00,	3,400.00,		
Titanium dioxide - (13463-67-7)	10,000.00,	10,000.00,		6.82,
xylene - (1330-20-7)	4,299.00,	1,548.00,		20.00,

( )		
( )		
( )		
/	2	
/	2	
	1	
(1 )	3	
( )		

12.

12.1.

Dangerous Preparations Directive 1999/45/EC

가  
.( 3 )



	96 hr LC50 mg/l	49 hr EC50 mg/l	ErC50 mg/l
xylene - (1330-20-7)	3.30, Oncorhynchus mykiss	8.50, Palaemonetes pugio	100.00 (72 hr), Chlorococcales
Epoxy resin (av.mol.wt.<700) - (25068-38-6)	3.10, Pimephales promelas	1.40, Daphnia magna	
Cyclohexanol, 4,4(1-methylethylidene)bis-, polymer with (chloromethyl)oxirane - (30583-72-3)			
Titanium dioxide - (13463-67-7)	1,000.00, Fundulus heteroclitus	5.50, Daphnia magna	5.83 (72 hr), Pseudokirchneriella subcapitata
Ethylbenzene - (100-41-4)	4.20, Oncorhynchus mykiss	2.93, Daphnia magna	3.60 (96 hr), Pseudokirchneriella subcapitata
Propylene glycol mono methyl ether - (107-98-2)	1,000.00, Oncorhynchus mykiss	500.00, Daphnia magna	1,000.00 (96 hr), Selenastrum capricornutum
Solvent naphtha (petroleum), light aromatic - (64742-95-6)	9.22, Oncorhynchus mykiss	6.14, Daphnia magna	19.00 (72 hr), Selenastrum capricornutum
1-Methoxy-2-propyl acetate - (108-65-6)	100.00, Salmo gairdneri	500.00, Daphnia magna	
polyamide dispersion - (55349-01-4)			

12.2.

가 .

12.3.

12.4.

12.5. , 가

PBT ( , ) vPvB ( , ) .

12.6.

13.

13.1.

가

..

14.

14.1.

1263

14.2.

14.3.

1263, , 3, III, 3[Y]

IMDG

Class/Div. 3

EmS

F-E,S-E

ICAO/IATA 3

14.4. III

14.5.

:

IMDG :

14.6. 가 가

14.7. MARPOL73/78 Annex II IBC Code .

15.

4 , 2 , III

MSDS 8 .

Ethylbenzene (0000100-41-4)  
Talc (0014807-96-6)  
Titanium dioxide (0013463-67-7)

(CMR):

carbon black (0001333-86-4)  
Ethylbenzene (0000100-41-4)  
Titanium dioxide (0013463-67-7)

:

Ethylbenzene (0000100-41-4)  
Titanium dioxide (0013463-67-7)  
xylene (0001330-20-7)

:

Ethylbenzene (0000100-41-4)  
xylene (0001330-20-7)

가 :

( )

:

( )

:

( )

:

( )

**Group I:**

( )

**Group II:**

Barium Sulphate (0007727-43-7)

Epoxy resin (av.mol.wt.<700) (0025068-38-6)

Ethylbenzene (0000100-41-4)

xylene (0001330-20-7)

:

( )

( )

:

Talc (0014807-96-6)

**16.**

: 10/25/2014

: 4

: 03/24/2006

MSDS KOSHA, NITE, ESIS, NLM, SIDS, IPCS, NCIS

SDS

Section 3

Phrases

H225

H226

H304

H312

H315

H317

H319

H332

H335

H336

H372

H373

H411

**This SDS is valid for 5 years from the revised date on page 1.**



Akzo Nobel

가