

ECB000/M INTERGARD 740 WHITE MATT

4

10/25/14

1.

1.1. INTERGARD 740 WHITE MATT
ECB000/M

1.2.

1.3.

626-6

(8-6)

1.4.

055-632-6286(),055 586 2310()

055 587 6276()

055 586 2310()

055 586 2310()

2.

2.1.

3; H226

/

2;H315

/

2;H319

1;H317

3;H412

2.2.

11 , 12



H315
H317
H319
H412

[]:

P210 / / /
P260 / /
P261 / /가 / / /
P262 , ,
P264
P272
P273
P280 / / /

[]:

P301+310 : /
P302+352 :
P303+361+353 () :
/
P305+351+338 가 : .가

P321 ().

P331

P333+313 /

P337 :

P362

P363

P370 :

P378 , , ,

[]:

P403+233 가

[]:

P501 ()

2.3.

PBT (,) vPvB (,)

3.

/	%	GHS	
EPOXY RESIN CAS No: 0025036-25-3	20-30	/ 2;H319 / 2;H315, 1;H317	[1]
Titanium dioxide CAS No: 0013463-67-7	10-20		[1][2]
xylene CAS No: 0001330-20-7	10-20	3; H226 - 4;H312 - 4;H332 / 2;H315 / 2A;H319 -1 ;H336 - 1;H372	[1][2]
Propylene glycol mono methyl ether	10-20		[1][2]

CAS No: 0000107-98-2		3; H226 -1 ;H336	
Fatty Acids, C18-unsaturated, Polymers with Bisphenol A and epichlorohydrin CAS No: 0067989-52-0	2.5-5	/ 2;H319 -1 ;H335 1;H317 - 2;H411	[1]
Solvent naphtha (petroleum), light aromatic CAS No: 0064742-95-6	2.5-5	1;H304	[1]
Ethylbenzene CAS No: 0000100-41-4	2.5-5	2;H225 - 4;H332 - 3;H373 1;H304 / 2;H315 / 2;H319 -1 ;H335	[1][2]
Amorphous fumed silica CAS No: 0112945-52-5	1-2.5		[1]
Amorphous Silica CAS No: 0007631-86-9	1-2.5		[1]
polyamide dispersion CAS No: 0055349-01-4	<1	1;H317 - 4;H413	[1]
	10-20	---	---

- 1)
- 2) 가
- 3) PBT vPvB
16

4.

4.1.

가

가

10

4.2. 가 /

4.3.

5.

5.1.

;

Note; 가

가

5.2.

가

:

5.3.

가

가

6.

6.1.

가 가 가 가

6.2.

가

6.3.

.8

, , 가 . (13 .)

가

가

, 가 , 가

7.

7.1.

가

, 가 (LEL) (OEL)

가

가

가 (LEL)

(OEL)

7.2. ()

, 8

가

가

61

가

1

7.3. Specific end use(s)

가

, 가

. 3

Hot surfaces, Sparks,

가

(60%

,)

8.

8.1.

(OEL)

(ACGIH)

(ACGIH)

ppm

mg/m³

ppm

mg/m³

Barium Sulphate

2

10

Ethylbenzene

125

545

100

435

Propylene glycol mono methyl ether

150

540

100

360

Talc

2

Titanium dioxide

10

xylene 150 655 100 434

(P) (Peak exposure limit)
(R)
(Sk)
(Sen)
(Cat 1)
(Cat 2) 가
(Cat 3)

DNEL/PNEC

8.2.

가

가

(visor)

(overall)

가

가

가

가

9.

pH

/ (°C)

(°C)

65

25

(= 1)

(,)

/

: 1.1 (xylene)

: 6.6 (xylene)

(Pa)

1.36

n- / (Log Kow)

9.2.

10.

10.1.

10.2.

.(Section 7)

가

10.3.

가

10.4.

.(7 .)

10.5.

10.6.

가

11.

(OEL)

가

가

가

2

LD50,
mg/kg

LD50,
mg/kg

LD50,

/

LD50,

			mg/L/4hr	mg/L/4hr
Amorphous fumed silica - (112945-52-5)	3,160.00,			
Amorphous Silica - (7631-86-9)	5,110.00,	5,000.00,		
EPOXY RESIN - (25036-25-3)				
Ethylbenzene - (100-41-4)	3,500.00,	15,433.00,	17.20,	
Fatty Acids, C18-unsaturated, Polymers with Bisphenol A and epichlorohydrin - (67989-52-0)				
polyamide dispersion - (55349-01-4)				
Propylene glycol mono methyl ether - (107-98-2)	5,000.00,	13,000.00,		
Solvent naphtha (petroleum), light aromatic - (64742-95-6)	6,800.00,	3,400.00,		
Titanium dioxide - (13463-67-7)	10,000.00,	10,000.00,		6.82,
xylene - (1330-20-7)	4,299.00,	1,548.00,		20.00,

()		
()		
()		
/	2	
/	2	
	1	
(1)		
()		

12.

12.1. Dangerous Preparations Directive 1999/45/EC 가 .(3 .)

가

	96 hr LC50 mg/l ,	49 hr EC50 mg/l ,	ErC50 mg/l ,
EPOXY RESIN - (25036-25-3)			
Titanium dioxide - (13463-67-7)	1,000.00, Fundulus heteroclitus	5.50, Daphnia magna	5.83 (72 hr), Pseudokirchneriella subcapitata
xylene - (1330-20-7)	3.30, Oncorhynchus mykiss	8.50, Palaemonetes	100.00 (72 hr), Chlorococcales

		pugio	
Propylene glycol mono methyl ether - (107-98-2)	1,000.00, Oncorhynchus mykiss	500.00, Daphnia magna	1,000.00 (96 hr), Selenastrum capricornutum
Fatty Acids, C18-unsaturated, Polymers with Bisphenol A and epichlorohydrin - (67989-52-0)			
Solvent naphtha (petroleum), light aromatic - (64742-95-6)	9.22, Oncorhynchus mykiss	6.14, Daphnia magna	19.00 (72 hr), Selenastrum capricornutum
Ethylbenzene - (100-41-4)	4.20, Oncorhynchus mykiss	2.93, Daphnia magna	3.60 (96 hr), Pseudokirchneriella subcapitata
Amorphous fumed silica - (112945-52-5)			
Amorphous Silica - (7631-86-9)	10,000.00, Danio rerio	10,000.00, Daphnia magna	10,000.00 (72 hr), Scenedesmus subspicatus
polyamide dispersion - (55349-01-4)			

12.2.

가 .

12.3.

12.4.

12.5. , , 가

PBT (,) vPvB (,) .

12.6.

13.

13.1.

가

14.

14.1. 1263

14.2.

14.3.

1263, , 3, III, 3[Y]

IMDG Class/Div. 3

EmS F-E,S-E

ICAO/IATA 3

14.4. III

14.5.

:

IMDG

:

14.6. 가

가

14.7.

MARPOL73/78 Annex II IBC Code .

15.

4

, 2

,

III

MSDS 8 .

Ethylbenzene (0000100-41-4)

Talc (0014807-96-6)

Titanium dioxide (0013463-67-7)

(CMR):

ethanol (0000064-17-5)

Ethylbenzene (0000100-41-4)

Titanium dioxide (0013463-67-7)

:

Ethylbenzene (0000100-41-4)

Titanium dioxide (0013463-67-7)

xylene (0001330-20-7)

:

Ethylbenzene (0000100-41-4)

xylene (0001330-20-7)

가

:

()

:

()

:

()

:

()

Group I:

()

Group II:

Barium Sulphate (0007727-43-7)

Ethylbenzene (0000100-41-4)

xylene (0001330-20-7)

()

()

Talc (0014807-96-6)

16.

: 10/25/2014

: 4

: 10/29/2010

MSDS KOSHA, NITE, ESIS, NLM, SIDS, IPCS, NCIS

SDS

Section 3 Phrases

H225

H226

H304

H312

H315

H317

H319

H332

H335

H336

H372

H373

H411

This SDS is valid for 5 years from the revised date on page 1.



Akzo Nobel

가